hygiprep

A simpler and easier bowel prep for colonoscopy

 Performed prior to your scheduled colonoscopy Replaces oral prep

The HygiPrep® is \$245 payable prior to performing the prep at the Hygieacare[®] Center. FSA, HRA, HSA eligible.

> For a consult and scheduling please call the **Hygieacare Center:** (734) 290-8437

5300 Elliott Dr. Ypsilanti, Michigan 48197



www.hygieacare.com



No need to run to the toilet



Get a good night's sleep* *No need to get up at night to take laxative



Achieve excellent outcomes

Over 20,000 patients chose Hygieacare® Prep

Excellent GI feedback on colon cleanliness 98% adequately clean

Excellent patient satisfaction 99% rate their experience "very good" or "good "



Perform HygiPrep® to effectively clean the colon - comfortably

hv**@1ea**care



1KM-059-

www.hygieacare.com

© Hygieacare, Inc. 2023. ALL RIGHTS RESERVED. No publication, distribution or reproduction of this document or any portion thereof is permitted without the written permission of Hygieacare, Inc.


HygiPrep®

A simpler and easier

bowel prep for

colonoscopy

Replaces oral prep





Safe • Easy • Convenient • Hygienic






hygiprep



Patient Instructions for HygiPrep® will be provided to you when scheduling your prep. A few days prior to your scheduled visit, you will receive a call to review the instructions.



Our trained technician will lead you to your private room, where you will be seated on the sanitized basin. A sterile disposable nozzle will be introduced into the rectum. A gentle stream of warm water will flow into the bowel, loosening stool.



Water continues to flow allowing you to comfortably and discreetly evacuate your colon. The trained technician will instruct you on completion of the procedure, which routinely takes less than 2 hours.

HygiPrep Patient feedback:



*Based on analysis of 2,468 patient written feedback for HygiPrep

The Hygieacare[®] System is an FDA cleared prescription medical device, approved only for colon cleansing, when medically indicated, such as before radiologic or endoscopic examinations. Possible side effects include abdominal cramping, nausea, vomiting, dizziness, and clinically insignificant rectal abrasions.